U.S.A. SUICIDE: 2003 OFFICIAL FINAL DATA

	Number	Per Day	Rate	% of Deaths	Group (Number of Sui.) Rate
Nation	31,484	86.3	10.8	1.3	White Male (22,830)19.5
Males	25,203	69.0	17.6	2.1	White Female (5,655) 4.7
Females	6,281	17.2	4.3	0.5	Nonwhite Male (2,373) 9.1
Whites	28,485	78.0	12.1	1.4	Nonwhite Female (626) 2.2
Nonwhites	2,999	8.2	5.5	0.9	Black Male (1,597) 8.8
Blacks	1,955	5.4	5.1	0.7	Black Female (358) 1.8
Elderly (65+ yrs.)	5,248	14.4	14.6	0.3	Hispanic (2,007) 5.0
Young (15-24 yrs.)	3,988	10.9	9.7	11.9	Native Americans (322)10.4
					Asian/Pacific Islanders (722). 5.5

Completions:

Black

7.0 6.7 6.5

a slight decline from 2002 was seen in 2003, but continues a pattern primarily of stability or slight decline in recent years

- Average of 1 person every 16.7 minutes killed themselves.
- Average of 1 old person every 1 hour 40.2 minutes killed themselves.
- Average of 1 young person every 2 hours 11.8 minutes killed themselves. (If the 250 suicides below age 15 are included, 1 young person every 2 hours 4 minutes)

• 11th ranking cause of death in U.S.— 3rd for young	>	> <u>Cause</u> 1	Number	Rate
• 4 male deaths by suicide for each female death by suicide.		All Causes	33,568	81.5
 Suicide ranks 11th as a cause of death; Homicide ranks 15th 		1-Accidents	15,272	37.1
Attempts (figures are estimates; no official U.S. national data are compiled):	1	2-Homicide	5,368	13.0
• 787,000 annual attempts in U.S. (using 25:1 ratio)	l	3-Suicide	3,988	9.7
• 25 attempts for every death by suicide for nation. 100-200:1 for young; 4:1 for elderly.		10-14 yrs	244	1.2
• 3 female attempts for each male attempt.		15-19 yrs	1487	7.3
		20-24 yrs	2501	12.1

Survivors (i.e., family members and friends of a loved one who died by suicide):

5.6 5.6 5.3 5.1

- Each suicide intimately affects at least 6 other people (estimate)
- Based on the 749,337 suicides from 1979 through 2003, estimated that the number of survivors of suicides in the U.S. is 4.5 million (1 of every 65 Americans in 2003); number grew by at least 188,904 in 2003.
- If there is a suicide every 16.7 minutes, then there are 6 new survivors every 16.7 minutes as well.

Suicide Methods		N	lumb	er	Rate I		Percent of Total				Number	Rate	Percer	t of Total			
Firearm suicides			16,90)7	5.8		53.7%			All but Firearms	14,577	5.0	46.	3%			
Suffocation/Hanging			6,63	35	2.3		21.1%			Poisoning 5,462 1.		1.9	17.3%				
Cut/pierce			57	71	0.2 1.8%			Drowning	339 0.1		1.	1.2%					
U.S.A. Suicide Rates 1994-2003 15 Leading Causes of Death in the U.																	
Group/	Group/ (Rates per 100,000 population) Group/							II	(total of 2,448,288 deaths; 841.9 rate)								
Age	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	Age	_	Rank & Cause of Dea	ath		Rate	Deaths
5-14	0.9	0.9	0.8	0.8	0.8	0.6	0.8	0.7	0.6	0.6	5-14	Ш	1 Diseases of heart (he	art disease)		235.6	685,089
15-24	13.8	13.3	12.0	11.4	11.1	10.3	10.4	9.9	9.9	9.7	15-24	Ш	2 Malignant neoplasms (cancer)			191.5	556,902
25-34	15.4	15.4	14.5	14.3	13.8	13.5	12.8	12.8	12.6	12.7	25-34	Ш	3 Cerebrovascular dise	ases (stroke)		54.2	157,689
35-44	15.3	15.2	15.5	15.3	15.4	14.4	14.6	14.7	15.3	14.9	35-44	Ш	4 Chronic lower respir	atory diseases		43.5	126,382
45-54	14.4	14.6	14.9	14.7	14.8	14.2	14.6	15.2	15.7	15.9	45-54	Ш	5 Accidents (unintention	onal injuries)		37.6	109,277
55-64	13.4	13.3	13.7	13.5	13.1	12.4	12.3	13.1	13.6	13.8	55-64	Ш	6 Diabetes mellitus (di	abetes)		25.5	74,219
65-74	15.3	15.8	15.0	14.4	14.1	13.6	12.6	13.3	13.5	12.7	65-74	\parallel	7 Influenza & pneumo	nia		22.4	65,163
75-84	21.3	20.7	20.0	19.3	19.7	18.3	17.7	17.4	17.7	16.4	75-84	Ш	8 Alzheimer's disease			21.8	63,457
85+	23.0	21.6	20.2	20.8	21.0	19.2	19.4	17.5	18.0	16.9	85+	Ш	9 Nephritis, nephrosis	(kidney disease)		14.6	42,453
65+	18.1	18.1	17.3	16.8	16.9	15.9	15.3	15.3	15.6	14.6	65+	Ш	10 Septicemia			11.7	34,069
Total	12.0	11.9	11.6	11.4	11.3	10.7	10.7	10.8	11.0	10.8	Total	- 11	11 Suicide [Intentiona	l Self-Harm]		10.8	31,484
Men	19.8	19.8	19.3	18.7	18.6	17.6	17.5	17.6	17.9	17.6	Men	Ш	12 Chronic liver disease	and cirrhosis		9.5	27,503
Women	4.5	4.4	4.4	4.4	4.4	4.1	4.1	4.1	4.3	4.3	Women	Ш	13 Essential hypertension	on and renal dise	ase	7.5	21,940
White	12.9	12.9	12.7	12.4	12.4	11.7	11.7	11.9	12.2	12.1	White	\parallel	14 Parkinson's disease			6.2	17,997
Nonwh	7.2	6.9	6.7	6.5	6.2	6.0	5.9	5.6	5.5	5.5	NonWh		15 Homicide [Assault]			6.1	17,732

Old made up 12.4% of 2003 population but represented 16.7% of the suicides. Young were 14.2% of 2003 population and comprised 12.7% of the suicides.

5.1 Black | - All other causes (Residual)

143.4

416,932

Official data source: CDC's WISQARS website and "Fatal Injury Reports": http://www.cdc.gov/ncipc/wisqars/; obtained 7 February 2006.

suicide rate = number of suicides by group Suicide Data Page: 2003

**Topopulation of group X 100,000

Prepared for AAS by John L. McIntosh, Ph.D.